



# **PUBLIC HEALTH**

**ALWAYS WORKING FOR A SAFER AND  
HEALTHIER WASHINGTON**

## **Rule Proposal to WAC 246-105-040, National Immunization Guidelines**

Washington State Board of Health  
June 13, 2012

Janna Bardi, Director, Office of Immunization and Child Profile

## Summary of Proposed Rule

Updates the reference to the Advisory Committee on Immunization Practices' (ACIP) national guidelines from the 2010 version to the 2012 version.

- Reflects changes to Tdap vaccine requirements in the board's Implementation Plan adopted on March 14, 2012.
- No changes to any other required vaccinations.

## Updated ACIP National Guidelines, Tdap Vaccine

2010	2012
Routinely recommended at 11 years old	No change
Five-year interval since a child's last DTaP or Td dose	Administer Tdap regardless of any interval
Undervaccinated children 7-10 years of age should receive <b>Td</b> vaccine	Undervaccinated children 7-10 years of age should receive <b>Tdap</b> vaccine

## Updated ACIP National Guidelines, Tdap Vaccine

- Providers should not delay Tdap; give it sooner to undervaccinated children, if needed
- No additional Tdap doses are established (one and done)
- ACIP's reasons for Tdap updates:
  - Additional data available for Tdap vaccination among 7-10 year-olds
  - Persistence of pertussis outbreaks in the U.S.
  - Protecting under-immunized kids against pertussis earlier prevents the chance of disease and its spread to babies.

**PUBLIC HEALTH**  
ALWAYS WORKING FOR A SAFER AND  
**HEALTHIER WASHINGTON**

Questions?

# Vaccine Requirements and Guidelines

**Recommended Immunization Schedule for Persons Aged 0 Through 6 Years -- United States, 2012 \***

Vaccine ▼	Age ►	Birth	1 month	2 months	4 months	6 months	9 months	12 months	15 months	18 months	19–23 months	2–3 years	4–6 years	
Hepatitis B <sup>1</sup>		Hep B	HepB			HepB								Range of recommended ages for all children
Rotavirus <sup>2</sup>				RV	RV	RV <sup>2</sup>								
Diphtheria, tetanus, pertussis <sup>3</sup>				DTaP	DTaP	DTaP	see footnote <sup>9</sup>	DTaP					DTaP	Range of recommended ages for certain high-risk groups
<i>Haemophilus influenzae</i> type b <sup>4</sup>				Hib	Hib	Hib <sup>4</sup>		Hib						Range of recommended ages for certain high-risk groups
Pneumococcal <sup>5</sup>				PCV	PCV	PCV		PCV				PPSV		Range of recommended ages for certain high-risk groups
Inactivated poliovirus <sup>6</sup>				IPV	IPV	IPV						IPV		
Influenza <sup>7</sup>						Influenza (Yearly)								
Measles, mumps, rubella <sup>8</sup>								MMR		see footnote <sup>9</sup>			MMR	Range of recommended ages for all children and certain high-risk groups
Varicella <sup>9</sup>								Varicella		see footnote <sup>9</sup>			Varicella	
Hepatitis A <sup>10</sup>								Dose 1 <sup>10</sup>			HepA Series			
Meningococcal <sup>11</sup>							MCV4 — see footnote <sup>11</sup>							

\* 2012 ACIP Immunization Schedule. Board-required vaccines for school and child care are circled in red.

# Vaccine Requirements and Guidelines

**Recommended Immunization Schedule for Persons Aged 7 Through 18 Years** -- United States, 2012 \*

Vaccine ▼	Age ►	7–10 years	11–12 years	13–18 years	
Tetanus, diphtheria, pertussis <sup>1</sup>		1 dose (If Indicated)	1 dose	1 dose (If Indicated)	Range of recommended ages for all children
Human papillomavirus <sup>2</sup>		See footnote <sup>2</sup>	3 doses	Complete 3-dose series	
Meningococcal <sup>3</sup>		See footnote <sup>3</sup>	Dose 1	Booster at age 16 years	Range of recommended ages for catch-up immunization
Influenza <sup>4</sup>		Influenza (yearly)			
Pneumococcal <sup>5</sup>		See footnote <sup>5</sup>			Range of recommended ages for certain high-risk groups
Hepatitis A <sup>6</sup>		Complete 2-dose series			
Hepatitis B <sup>7</sup>		Complete 3-dose series			
Inactivated poliovirus <sup>8</sup>		Complete 3-dose series			
Measles, mumps, rubella <sup>9</sup>		Complete 2-dose series			
Varicella <sup>10</sup>		Complete 2-dose series			

\* 2012 ACIP Immunization Schedule. Board-required vaccines for school and child care are circled in red.